

The Fundamental Structure of Coronal Loops

Dr. Amy R. Winebarger (NASA MSFC)

H.P. Warren (NRL), J.W. Cirtain (NASA MSFC), K. Koboayashi (UAH), K. Korreck (SAO), L. Golub (SAO), S. Kuzin (LI), R.W. Walsh (UCLAN), C. DeForrest (SWRI), B. DePointieu (LMSAL), A. Title (LMSAL), M. Weber (SAO)

High Resolution Coronal Imager (Hi-C)



- Flown on a sounding rocket from White Sands on July 11, 2012
- Observed AR 11520 for ~ 5 minutes
- 193 Å channel
- 0.1'' pixels, ~0.25'' resolution
- 6.8' x 6.8' field of view
- 5.4 s cadence


Highest-resolution, fastest-cadence EUV images
taken of the corona.

What to take away

- Hi-C reveals that loops have sub-structure below what can be resolved with AIA.
- The strands appear to be twisted.
- We may need higher resolution yet...


Hi-C Field of View








AIA 193 Å 11-Jul-12 18:55:31.840

Evolving Loop




Hi-C resolves several strands wrapping around one another.

Evolving Loop




Evolving Loop




Shortly after Hi-C, an event occurred at the crossing.

Are we resolving the sub-structure?




EIS, Hi-C and AIA Rasters

← time







Isolate a structure



Is this structure resolved?


Calculate a density



$$n_e = 2-7 \times 10^{10} \text{ cm}^{-3}$$

$$n_e = 0.5-6 \times 10^{10} \text{ cm}^{-3}$$

Potentially resolved



To replicate Hi-C emission:

$$T = 1.5 \text{ MK}$$

Radius of structure = 435 km

$$n_e = 1 \times 10^{10} \text{ cm}^{-3}$$

Densities measured from EIS: $0.5\text{-}7 \times 10^{10} \text{ cm}^{-3}$

Hi-C: A New View of the Sun

Data will be released in January.